

Kai Mithoefer, MD

New England Center for Regenerative Orthopedics and Sports Medicine

Boston Sports and Shoulder Center

840 Winter Street, Waltham, MA 02451, (781) 890-2133

125 Parker Hill Avenue, Boston, MA 02120, (617) 264-1100

Postoperative Rehabilitation Guidelines

Meniscal Root Repair

The intent of this protocol is to provide the clinician with a guideline of the post-operative rehabilitation course of a patient that has undergone a meniscal root repair. It is not means intended to be a substitute for one's clinical decision making regarding the progression of a patient's post-operative course based on their physical exam/findings, individual progress, and/or the presence of post-operative complications. If a clinician requires assistance in the progression of a post-operative patient, they should consult with the referring Surgeon.

Progression to the next phase based on clinical criteria and/or time frames as appropriate.

Key factors in determining progression of rehabilitation after Meniscal repair include:

- Tissue quality at Time of repair
- Other pathology (ligamentous injury)

Phase 1- Maximum Protection - Weeks 1-4:

Goals:

- Diminish inflammation and swelling
- Restore ROM
- Reestablish quadriceps muscle activity

Stage 1: Immediate Postoperative Day 1 – Week 3

- Ice, compression, elevation
- Electrical muscle stimulation
- Brace for support with ROM limitation
- ROM 0-110

- Motion is limited for the first 7-21 days, depending on the development of scar tissue around the repair site. Gradual increase in flexion ROM is based on assessment of pain and site of repair (0-110 degrees)
- Patellar mobilization
- Scar tissue mobilization
- Passive ROM
- Exercises
 - Quadriceps isometrics
 - Hamstring isometrics (if posterior horn repair, no hamstring exercised for 6 weeks)
 - Hip abduction and adduction
- Weight-bearing: TDWB weeks 1+2, 50% weight bearing week 3+4. WBAT at weeks 4-6.
- Proprioception training with brace locked at 0 degrees

Stage 2: Weeks 4-6

- Progressive resistance exercises (PREs) 1-5 pounds
- Limited range knee extension (in range less likely to impinge or pull on repair)
- Toe raises
- Mini squats less (than 45 degrees flexion)
- Cycling (no resistance)
- Unloaded flexibility exercises

Phase 2: Moderate Protection Weeks 6-10

Criteria for progression to Phase 2:

- ROM 0-110 degrees
- No change in pain or effusion
- Quadriceps control (MMT 4/5)

Goals:

- Increased strength, power, endurance
- Normalize ROM of knee
- Prepare patients for advanced exercises

Exercises:

- Strength – Pre progression
- Flexibility exercises
- Lateral step-ups
- Mini squats

Endurance Program:

- Swimming (no frog kick), pool running-if available
- Cycling
- Stair machine

Coordination Program:

- Pool sprinting-if pool available

- Backward walking

Phase 3: Advanced Phase Weeks 11-15

Criteria for progression to Phase 3

- Full, pain free ROM
- No pain or tenderness
- Satisfactory clinical examination
- SLR without lag
- Gait without device, brace unlocked

Goals:

- Increase power and endurance
- Emphasize return to skill activities
- Prepare for return to activities

Exercises:

- Continue all exercises
- Plyometrics, pool program

Phase 4: Activity Progression Phase

Progression Criteria

- Full, pain free ROM
- Satisfactory clinical examination

Criteria for discharge from skilled therapy:

- Non-antalgic gait
- Pain free/full ROM
- LE strength at least 4/5
- Normal age appropriate balance and proprioception
- Resolved palpable edema

Goals:

- Prepare for return to full unrestricted activities

Exercises:

- Continue all exercises
- Advance plyometrics
- Agility program
- Sport-specific functional progression
- Independent with home program